

Use of Gödel Universe to Construct A New Zollfrei Metric with $R^2 \times S^1$ Topology

Moninder Singh Modgil ¹

Abstract

A new example of $(2 + 1)$ -dimensional Zollfrei metric, with the topology $R^2 \times S^1$, is presented. This metric is readily obtained from the celebrated $(3 + 1)$ - dimensional rotating Gödel universe $G_{3,1}$. This is because $G_{3,1}$ has the interesting property that, the light rays which are confined to move on the plane perpendicular to the rotation axis, return to their origin after a time period $T = \frac{2\pi}{\omega}[\sqrt{2} - 1]$ -where ω is the angular velocity of the universe. Hence by - the topological identification of pairs of points on the time coordinate, seperated by the time interval T . and dropping the flat x_3 coordinate - which is directed along the rotation axis; one obtains the $(2 + 1)$ -dimensional Zollfrei metric with the $R^2 \times S^1$ topology.

KEY WORDS: Gödel Universe, Zollfrei Metric, Closed Null Geodesics (CNCs), Closed Timelike Curves (CTCs), Periodic Time

¹ PhD in Physics, from Indian Institute of Technology, Kanpur, India, and B.Tech.(Hons.) in Aeronautical Engineering, from Indian Institute of Technology, Kharagpur, India.
email: moni_g4@yahoo.com

A Lorentzian manifold is said to possess Zollfrei metric, if all its null geodesics are closed [2]. Part of attraction for studying Zollfrei metrics in physical context comes from the Segal's cosmological model [4] which is based upon the $S^3 \times S^1$ topology of spacetime. Examples of manifolds with Zollfrei metric are - $S^n \times S^1$ and $P^n \times S^1$ - with commensurate radii of spatial and temporal factors. To our knowledge, all previously known examples of Zollfrei metric have compact topology for the spatial factor - i.e., S^n or P^n . Here, we construct a new example of Zollfrei metric in $(2+1)$ -dimensions, with the $R^2 \times S^1$ topology. This construction is based upon the Gödel universe [1]. We shall refer to this as the Gödel-Zollfrei metric, and denoted it by $G_{2,1}^{S^1}$.

The $(3+1)$ -D axisymmetric, homogenous, Gödel universe $G_{3,1}$ [1] with the topology $R^3 \times R^1$, has the line element -

$$ds^2 = a^2(dx_0^2 - dx_1^2 - dx_3^2 + \frac{e^{2x_1}}{2}dx_2^2 + 2e^{x_1}dx_0dx_2) \quad (1)$$

which satisfies the Einstein equations for a uniform matter density,

$$\rho = (8\pi Ga^2)^{-1}, \quad (2)$$

and a cosmological constant,

$$\Lambda = -4\pi G\rho, \quad (3)$$

where, G is Newton's gravitational constant. The universe rotates about the x_3 axis, with the angular velocity,

$$\omega = 2(\pi G\rho)^{1/2} = (\sqrt{2}a)^{-1} \quad (4)$$

Lets denote the curved space defined by the three coordinates (x_0, x_1, x_2) as $G_{2,1}$. Now $G_{3,1}$ can be regarded as the product of $G_{2,1}$ and the flat x_3 coordinate (which has the topology R^1), i.e.,

$$G_{3,1} = G_{2,1} \times R^1 \quad (5)$$

Pfarr [3] worked out geodesic and non-geodesic trajectories for $G_{2,1}$. For particles confined to move on the (x_1x_2) plane - i.e., the plane perpendicular to the rotation axis x_3 , he showed that the geodesically moving

particles, move on circles and consequently return to the point of their origin. In particular, the light rays move on the largest circles (null geodesics), and reach out to a maximal distance r_H , from the point of their origin,

$$r_H = \frac{2}{\omega} \ln(\sqrt{2} + 1) \quad (6)$$

The circle defined by r_H , may be termed as the Gödel horizon. The circles of radius $r = r_H$ are Closed Null Curves (CNCs), while circles of radius $r > r_H$ are Closed Timelike Curves (CTCs). The time taken by the light rays to return to the point of their origin is [3] -

$$T = \frac{2\pi}{\omega} [\sqrt{2} - 1] \quad (7)$$

Accordingly if in $G_{3,1}$ -

1. One drops the flat x_3 coordinate, which is directed along the rotation axis, and
2. Compactifies the time coordinate x_0 , to S^1 , and
3. Chooses the time period of S^1 time factor equal to $T = \frac{2\pi}{\omega} [\sqrt{2} - 1]$;

one obtains this new example of $(2 + 1)$ -dimensional Zollfrei metric with the $R^2 \times S^1$ topology.

Gödel [1] gave a list of nine interesting properties of $G_{3,1}$. Remarkably, in the fifth property, he considered the possibility of both an open R^1 and a closed S^1 time coordinate.

References

- [1] Gödel, K.: *Rev. Mod. Phys.* **21**, (1949), 447.
- [2] Guillemin, V.: *Cosmology in (2+1)-dimensions, cyclic models and deformations of $M_{2,1}$* , Princeton University Press, (1989).
- [3] Pfarr, J.: *Gen Rel. and Grav.*, **13**, 1073, (1981).
- [4] Segal I.E. : *Mathematical Cosmology and Extra-galactic Astronomy*, Cambridge University Press (1976).